



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Inspection at El Paso, Tex.

Acting Assistant Surgeon Alexander reports as follows for the week ended August 20, 1904:

Mexican Central passengers inspected, 237; Mexican immigrants inspected, 64; inspection private Pullman passengers and crew, 18; disinfection of soiled linen imported for laundry work, 317 pieces; vaccination of children of immigrants, 4.

Inspection at Laredo, Tex.

Acting Assistant Surgeon Hamilton reports, through Passed Assistant Surgeon Richardson, as follows:

Week ended August 20, 1904:

Passenger trains from Mexico inspected, 14; persons on trains from Mexico inspected, 411; immigrants on trains from Mexico inspected, 18; immigrants vaccinated upon entry, 2; Pullman coaches disinfected, 14; persons detained for noncompliance with State quarantine regulations, 7; persons inspected and entered via international footbridge, 3,372; persons refused entry, presenting at international footbridge, 28.

Statistical reports of States and cities of the United States—Yearly and monthly.

CALIFORNIA.—Reports to the State board of health for the month of July, 1904, from 34 localities, having an aggregate population of 1,081,750, show a total of 1,241 deaths, including 107 from zymotic diseases and 182 from tuberculosis.

Sacramento.—Month of July, 1904. Estimated population, 30,000. Total number of deaths, 36, including enteric fever 1 and 5 from phthisis pulmonalis.

COLORADO.—Denver.—August 1 to 17, 1904. Estimated population, 190,000. Total number of deaths, 96, including diphtheria, 3; enteric fever, 1; whooping cough, 1, and 16 from phthisis pulmonalis.

CONNECTICUT.—Reports to the State board of health for the month of July, 1904, from 167 towns, having an aggregate estimated population of 955,496, show a total of 1,405 deaths, including diphtheria 9, enteric fever 12, measles 3, scarlet fever 10, whooping cough 6, and 100 from phthisis pulmonalis.

Bridgeport.—Month of July, 1904. Estimated population, 82,128. Total number of deaths, 135; including diphtheria, 2; enteric fever, 1; scarlet fever, 3; and 14 from tuberculosis.

ILLINOIS.—Peoria.—Month of July, 1904. Census population, 56,100. Total number of deaths, 74; including diphtheria, 1; and 6 from tuberculosis.

Rockford.—Month of July, 1904. Estimated population, 37,000. Total number of deaths, 29, including 5 from tuberculosis.

IOWA—*Dubuque*.—Three weeks ended August 20, 1904. Estimated population, 40,000. Total number of deaths, 25, including 4 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended August 20, 1904, from 78 observers, indicate that bronchitis, cholera infantum, dysentery, intermittent fever, erysipelas, and measles were less than usually prevalent, and smallpox, pleuritis, scarlet fever, diphtheria, and puerperal fever were more than usually prevalent.

Meningitis was reported present at 2, pneumonia at 8, whooping cough at 14, measles at 15, diphtheria at 23, scarlet fever at 36, smallpox at 58, enteric fever at 63, and phthisis pulmonalis at 280 places.

MINNESOTA—*Duluth*.—Month of July, 1904. Estimated population, 70,000. Total number of deaths, 56, including diphtheria 1, and 7 from tuberculosis.

NEW HAMPSHIRE—*Manchester*.—Month of July, 1904. Estimated population, 62,000. Total number of deaths, 101, including enteric fever 1, whooping cough 1, and 5 from tuberculosis.

NEW JERSEY—*Hudson County*.—Month of July, 1904. Estimated population, 433,573. Total number of deaths, 994, including diphtheria 21, enteric fever 5, measles 8, scarlet fever 8, whooping cough 6, and 89 from phthisis pulmonalis.

OHIO—*Cleveland*.—Month of July, 1904. Estimated population, 430,000. Total number of deaths, 562, including diphtheria 1, enteric fever 5, measles 11, and 52 from tuberculosis.

East Liverpool.—Month of July, 1904. Estimated population, 20,000. Total number of deaths, 11, including 1 from whooping cough.

TENNESSEE—*Knoxville*.—Month of July, 1904. Estimated population, 40,000; white, 30,000, colored, 10,000. Total number of deaths, 59; white, 43; colored, 16, including enteric fever 2, and 13 from tuberculosis.

Nashville.—Month of July, 1904. Estimated population, 82,947; white, 52,061; colored, 30,886. Total number of deaths, 202; white, 109; colored, 93, including enteric fever 7, measles 1, and 34 from tuberculosis.

UTAH.—Bulletin of the State board of health for July, 1904: Reports received from 91 local health officers in 26 counties, 42 of whom reported their communities free from all of the diseases required to be reported, show scarlet fever present in 7 localities, with no deaths; smallpox present in 6 localities, with no deaths; diphtheria present in 10 localities, with 6 deaths; enteric fever present in 15 localities, with 2 deaths; whooping cough present in 19 localities, with 2 deaths; measles present in 12 localities, with no deaths; pneumonia present in 9 localities, with 12 deaths; tuberculosis present in 8 localities, with 5 deaths; total, 27 deaths from contagious diseases.